



The Scientific Temper

VOL-VIII, NO.1&2; JANUARY-JULY, 2017

ISSN 0976 8653, E ISSN 2231 6396

UGC SR NO 2535; JR NO. 47226

e-mail:letmepublish@rediffmail.com

Web: www.scientifictemper.com

ECOLOGICAL ENGINEERING OF MICROALGAE FOR ENHANCED ENERGY PRODUCTION

Vijetna Singh* and Alka Thakur**

Department of Electrical Engineering,

Sri Satya Sai University of Technology and Medical Sciences,
Sehore (Madhya Pradesh), INDIA

ABSTRACT

Microalgae are considered as potential source of renewable energy generation. Algal biomass is used as key feedstock for renewable energy generation using algal resources. Accumulation of biomass mostly depends on the growth rate, which is usually regulated by pH and salinity. Present study was conducted to study the effect of variation in pH and saline ranges on growth, biomass and lipid content of a freshwater microalga *Chlorella vulgaris*. The growth, biomass and lipid content showed a significant difference due to the variation in pH and saline range. The growth and lipid content was maximum at pH 7.0 with compared to 6, 8 and 9. On the other hand an increase of salinity from 0.1M to 0.25 M caused decreased growth rate, whereas the lipid content showed an increasing trend.

Keywords: Biomass, *Chlorella vulgaris*, Growth, Lipids, Microalgae, pH, Salinity.

INTRODUCTION

Human civilization is facing several inter connected challenges such as energy supply, food security, health, among them uninterrupted energy (electricity) supply is the major cause of concern due its imp in every aspect of human life. Global energy particularly electricity generation is mainly dependent on non-renewable resources which are depleting gradually. On the other hand global energy

requirement is increasing at an alarming rate to meet the continuously growing demand of energy (electricity) supply of the world, alternate and renewable sources of energy or electricity production are being considered. Solar and wind energy are key sources of renewable power (electricity production) however a more efficient and eco-friendly way to generate energy which is based on biological resources is still in fancy state

few countries are however successfully harnessing biological energy but extensive research are still warranted to standardized and effectively biological resources of electricity production.

The continued use of fossil fuel has now recognized as unsustainable because of depleting supplies and accumulation of CO₂ and other Greenhouse gases (Patil *et al*; 2008). The combustion of fossil fuel leads to 73% of CO₂ production (Verma *et al*; 2009), which adversely affect the atmosphere. Therefore, the demand of biofuels is the necessity in today's scenario as it is made from non-toxic, biodegradable and renewable resources and provides environmental benefits (Gouveia and Oliveira; 2008). The use of biofuel leads to decrease in harmful emission of carbon monoxide, hydrocarbons, particulate matter and to the elimination of SO_x emissions with consequent decrease in Green House effect (Gouveia and Oliveira; 2008). The type of alternative sources of energy includes, hydroelectricity, energy generated from nuclear, wave and wind power and biological material (Scragg *et al*; 2003).

Among alternative resources of energy production, biological resources have received considerable attention. Many countries are harnessing the potential of biological resources for energy production. *Jatropha* and other similar seed plants have been widely recognized as sources of bioenergy, but cultivation of such plants may compromise the agricultural land. Therefore, scientists are considering algae as next generation sources of biological energy (Williams; 2007).

Production of biodiesel using microalgal biomass appears to be viable alternative. Microalgae are diverse group of prokaryotic and eukaryotic photosynthetic organisms that grow rapidly due to their simple structure (Yanqun Li *et al*; 2008). Microalgae are unicellular photosynthetic organisms, which uses sunlight and CO₂ for production of biomass they are considered having more photosynthetic efficiency than plants (Converti *et al*; 2009, Benemann; 1997 and Miao and Wu; 2006).

Rapid growth, large biomass and greater accumulation of lipid content in the algal cell are

the important criteria for considering the algal species as potential sources of energy. In this context, the present study is aimed to enhanced the growth rate, biomass production and lipid accumulation in a common freshwater alga *Chlorella vulgaris* through ecological engineering e. g. manipulation of external pH and salinity.

2. MATERIAL AND METHODS

2.1. Test Organism and growth conditions

A locally isolated freshwater green alga *Chlorella vulgaris* was used as test organism and cultivated in BG11 medium at pH 7.0 in a culture room at 25 ± 2° C under a photoperiod of 12:12 hour at light intensity of 75 μmol photon m⁻² s⁻¹ PAR.

The composition of the BG 11 medium was as follows: 1.5g NaNO₃, 0.04g K₂HPO₄, 0.075g MgSO₄.7H₂O, 0.036g CaCl₂.2H₂O, 0.006g Citric Acid, 0.006g Ferric Ammonium Citrate, 0.001g EDTA (Disodium magnesium salt) and 0.02g Na₂CO₃, and 2.286mg H₃BO₃, 1.81mg MnCl₂.4H₂O, 0.222mg ZnSO₄.7H₂O, 0.039mg Na₂MoO₄.2H₂O, 0.079mg CuSO₄.5H₂O, 0.0494 mg Co(NO₃)₂.6H₂O. pH was set between 7.0-7.5. The cultures were hand shaken two or three times daily to avoid sticking. This was referred to as control culture.

2.2. Optimization of medium on different pH and Salinity ranges

The one set of triplicates of cultures for evaluation were grown in 150 ml Erlenmeyer flasks containing 50 ml BG11 medium on different pH ranges (6.0–9.0 at an interval of 1.0), and another set of triplicates for different salinity ranges (0.1 M - 0.25 M at an interval of 0.05, for saline cultures the pH was set at 7.0) in a culture room at 25 ± 2° C under a photoperiod of 14:10 h at light intensity of 75 μmol photon m⁻² s⁻¹ PAR without sparging with air or CO₂.

Growth measurement

The growth of the algae was determined by measuring absorbance at 663 nm algal growth was measured on regular intervals (7, 14, 21 and 28 day respectively) using a spectrophotometer.

Measurement of dry weight

Dry weight (dcw) was determined gravimetrically. A known volume of algal culture was centrifuged at 5000 rpm for 15 min and the harvested biomass was dried at 80°C to constant weight and the weight of the dried biomass was recorded manually.

3. Extraction and Estimation of lipid from algal biomass

Extraction of lipid was performed following the protocol of Bligh and Dyer (1959). To a 15 ml glass vial containing a known amount of algal biomass, 2 ml methanol and 1 ml chloroform were added and kept for 24 h at room temperature. The mixture was agitated in a vortex for 2 min, and 1 ml of chloroform was again added and the mixture shaken vigorously for 1 min; 1.8 ml of distilled water was added and the mixture was agitated in a vortex again for 2 min. The layers were separated by centrifugation for 10 min at 2000 rpm. The lower layer was filtered through Whatman No. 1 filter paper into a previously weighed clean vial (W1). Evaporation was carried on in a water bath and the residue was further dried at 104°C for 30 min. The weight of the vial was again recorded (W2). Lipid content was calculated by subtracting W1 from W2, and was expressed as % dcw.

3.1. Statistical Analysis

All the experiments were performed in triplicates, the results were analyzed statistically by SPSS for two-way ANOVA.

4. RESULT AND DISCUSSION

4.1. Effect of varied pH on Growth and Biomass




Figure 1:

Growth Absorbance of *Chlorella vulgaris* with reference to the days of incubation

The Growth curve of *C. vulgaris* is shown in figure 1, the maximum algal density was statistically significant ($P < 0.05$) in control culture (pH 7) compared to pH 6, 8 and 9. The algal density was increased in all pH with passage of time. But control culture with pH 7 showed approximately, 40.0, 53.0 and 56% increased growth compared to pH 6.0; 8.0 and 9.0, respectively, on day 21. In control culture (pH 7) algal density exhibited increase of 83% from 7day to 21 day.




Figure 2: Biomass concentration of *Chlorella vulgaris* at variable pH ranges with reference to the days of incubation

Experiments were performed in triplicates under optimized conditions to determine the biomass yield, presented in figure 2, the most significant increase in biomass yield to 87.0 % was obtained in control culture on day 21 compared to pH 6, 8 and 9. pH 7 and pH 6 showed difference of 53.7%, whereas pH 8 and pH 9 showed 57.4% and 62.8% difference respectively on day 7 in comparison to control culture. The biomass yield was continuously increasing till day 21st after that no further increase was observed.

4.2. Effect of varied pH on Lipid Accumulation




Figure 3: Relationship between pH of the culture medium of *Chlorella vulgaris* and lipid accumulation with reference to the days of incubation

Experiments conducted at varied pH to study lipid accumulation in *C. vulgaris* resulted in increase/decrease in lipid yield with respect to control (pH 7) shown in Figure 3. pH 6 showed decrease in lipid pool compared to pH 7. Lipid yield showed increase of 124 folds and 178 folds respectively, in pH 8 and pH 9 on day 21 and decline was observed on day 28.

4.3. Effect of Salinity

4.3.1. Effect of Salinity on Growth




Figure 4: Growth of *Chlorella vulgaris* in medium supplemented with variable salinity ranges

The growth curve of *C. vulgaris* with variable salinity is shown in figure 4. Statistical analysis conclude that growth differs significantly ($P < 0.05$) with variable salinity. Tukey's post-hoc analysis suggests that control culture showed maximum growth and it decreased as salinity in medium was increased from 0.1M to 0.25M. Approximately, 46fold increase was observed on day 21 in the control culture compared to zero day. The increasing trend of growth was observed from zero day to 21st day in all concentrations of salinity after, which decline was observed on 28th day. Lowest growth was observed in culture with 0.25M salinity, an increase of 16.9 fold was observed after 21 days compared to zero day.




Fig. 5: Biomass of *Chlorella vulgaris* in medium supplemented with variable salinity ranges

The biomass also showed similar trend as noticed growth *i.e* highest biomass was observed in control culture and minimum in culture growing in 0.25 M saline medium. The increasing trend was observed from zero days to 28th day. Highest 87 fold increase was observed in control culture on day 28th. 0.25 M saline culture showed only 52 fold increase with respect to zero days.

4.3.2. Effect of Salinity on Lipid Content




Fig. 6: Lipid accumulation in *Chlorella vulgaris* at different concentration of NaCl

The lipid accumulation in *C. vulgaris* at different saline ranges was studied in triplicates under optimized conditions and it differed significantly ($P < 0.05$). The highest lipid yield was observed in culture growing in 0.25 M saline medium and lowest was observed in control culture (0.0M saline medium).

The yield increased from zero days to 28th day. On 2th day 91 fold increase was observed in 0.25 M saline culture compared to 81 fold, 72 fold, 66 fold and 42 fold increase in 0.2M, 0.15 M, 0.1 M and control culture i.e 0.0 M, respectively (Fig. 6).

The experiments conducted to study growth of *C. vulgaris* with variable pH showed that maximum growth occurs at control pH 7, whereas increasing pH decreased the algal density. It can be correlated to the fact that higher pH limits carbon availability from CO₂, hence algal growth is suppressed (Chen and Durbin; 1994, Azov; 1982 and Murthy *et al*; 2013). The biomass extracted from all variable pH ranges also showed highest biomass at control pH7. Since biomass is directly proportional to algal growth. Therefore, growth and biomass showed almost similar trend. The lipid accumulation at pH 9 was highest perhaps due to the fact that alkaline pH indirectly results in increased triglyceride accumulation (Guckert and Cooksey *et al* and

Murthy *et al*; 2013) alkaline pH indirectly results in increase in triglyceride accumulation.

Exposing algae to lower or higher salinity levels than their natural levels can change growth rate and alter composition. In our experiments the growth rate and biomass decreased with increasing NaCl concentration in the growth medium and hence highest growth was observed in 0.0M culture and lowest was observed in culture grown in 0.25 M salinity. This could be associated with the fact that *C.vulgaris* was unable to adapt at higher saline ranges. Cultivation with different salinity ranges although showed similar time course i.e growth increased from 0 to 21st day and declined on day 28. Higher lipid content was however observed in the medium with highest NaCl concentration i.e at 0.25 M.

ACKNOWLEDGEMENT:

I am grateful to the Head, Ms. Alka Thakur (Associate Professor, Department of Electrical Engineering) of SSSUTMS Sehore (M.P) for helping me in the various ways.

REFERENCES

1. Azov, Y. (1982). Effect of pH on inorganic carbon uptake in algal cultures. *Appl. Environ. Microbiol* 43, 1300–1306.
2. Benemann, J. (1997). Feasibility analysis of photobiological hydrogen production. *Int. J. Hydrog. Energy* 22, 979–987.
3. Bligh, E.G and Dyer, W.J. (1959). A rapid method of total lipid extraction and purification. *Can J BiochemPhysiol*37:911-917.
4. Chen, C.Y.; Durbin, E.G. (1994). Effects of pH on the growth and carbon uptake of marine phytoplankton. *Mar. Ecol.-Prog. Ser.* 109, 83–94.
5. Converti, A.; Casazza, A.A.; Ortiz, E.Y.; Perego, P.; del Borghi, M. (2009). Effect of temperature and nitrogen concentration on the growth and lipid content of *Nannochloropsisoculata* and *Chlorella vulgaris* for biodiesel production. *Chem. Eng. Process.* 48, 1146–1151.
6. Gouveia L and Oliveira A C (2008). Microalgae as raw material for biofuels production. *J IndMicrobiolBiotechnol* 36:269-274.
7. Guckert, J.B.; Cooksey, K.E. (1990). Triglyceride accumulation and fatty acid profile changes in *Chlorella* (Chlorophyta) during high pH induced cell cycle inhibition. *J. Phycol.* 26, 72–79.

8. Illman AM, Scragg AH, Shales SW (2000). Increase in *Chlorella* strains calorific values when grown in low nitrogen medium. *Enzyme Microb Technol* 27:631–635.
9. Khan, S.A.; Hussain, M.Z.; Prasad, S.; Banerjee, U. (2009). Prospects of biodiesel production from microalgae in India. *Renew. Sustain. Energy Rev.* 13, 2361–2372.
10. Mata T M, Martins AA and Nidia S C (2009). Microalgae for biodiesel production and other applications: A review. *Renewable and Sustainable Energy Reviews* 14, 217–232.
11. Miao, X and Wu, Q. (2006). Biodiesel production from heterotrophic microalgal oil, *Bioresource Technology*. 97,841-846.
12. Murthy G S, Ceballos R M and Juneja A (2013). Effect of Environmental Factors and Nutrient Availability on the Biochemical Composition of Algae for Biofuels Production: A Review. *Energies*,6, 4607-4638.
13. Ormerod, W. G., P. Freund, A. Smith, and J. Davison (2002). Ocean storage of CO₂, Greenhouse Gas R&D Programme, Int. Energy Agency, Cheltenham, U. K.
14. Patil, V., Tran, K. & Giselrod, H.R. (2008). Towards Sustainable Production of Biofuels from Microalgae. *International Journal of Molecular Science* 9: 1188-1195.
15. Rai, L.C; Mallick, N; Singh, J.B and Kumar, H.D. (1991). Physiological and biochemical characteristics of copper tolerant and wild type strain of *Anabaena doliolum* under copper stress. *J Plant Physiol* 138:68-74.
16. Scragg A.H, Morrison J and Shales S.W (2003) The use of a fuel containing *C. vulgaris* in a diesel engine. *Enzyme and microbial Technology* 33: 884-889.
17. Spolaore, P.; Joannis-Cassan, C.; Duran, E.; Isambert, A. (2006). Commercial applications of microalgae. *J. Biosci. Bioeng.* 101, 87–96.
18. Verma N M, Mehrotra S, Shukla A and Mishra B N (2009), Prospective of biodiesel production utilizing microalgae as the cell factories: A comprehensive discussion. *African Journal of Biotechnology* Vol. 9 (10), 1402-1411.
19. William T. Coyle (2007). The future of biofuels: A global perspective. *Amber waves*.
20. Yanqun L, Horsman M, Wang B, Wu N and Lan C Q (2008), Effects of nitrogen sources on cell growth and lipid accumulation of green alga *Neochloris oleoabundans*. *Applied Microbiology and Biotechnology* Volume 81, Issue 4, 629-636.

<http://www.scientifictemper.com/>